


**Risk
analysis
for plants
as pests
for *Ambrosia
trifida***



**Risk
analysis
for plants
as pests
for *Ambrosia
trifida***

Inter-American Institute for Cooperation on Agriculture (IICA), 2018



Risk analysis for plants as pests for *Ambrosia trifida* by IICA is published under license Creative Commons Attribution-ShareAlike 3.0 IGO (CC-BY-SA 3.0 IGO) (<http://creativecommons.org/licenses/by-sa/3.0/igo/>)

Based on a work at www.iica.int

IICA encourages the fair use of this document. Proper citation is requested.

This publication is available in electronic (PDF) format from the Institute's Web site: <http://www.iica.int>

Editorial coordination: Lourdes Fonalleras and Florencia Sanz

Translator: Alec McClay

Layout: Esteban Grille

Cover design: Esteban Grille

Digital printing

Risk analysis for plants as pests for *Ambrosia trifida* / Inter-American Institute for Cooperation on Agriculture, Comité Regional de Sanidad Vegetal del Cono Sur; Alec McClay. – Uruguay : IICA, 2018.

26 p.; A4 21 cm X 29,7 cm.

ISBN: 978-92-9248-811-6

Published also in Spanish and Portuguese

1. Asteraceae 2. *Ambrosia* 3. Phytosanitary measures
4. Pests of plants 5. Risk management 6. Pest monitoring 7. Weeds I. IICA II. COSAVE III. Title

AGRIS
H10

DEWEY
632.5

Montevideo, Uruguay - 2018

ACKNOWLEDGMENTS

The *Guidelines of procedures for risk assessment of plants as pests (weeds)* has been applied for the development of two case studies: *Hydrocotyle batrachium* and *Ambrosia trifida*. This products was a result of the component aimed to build technical capacity in the region to use a Pest Risk Analysis process with emphasis on the assessment of Plants as Pests (weeds) in the framework of STDF / PG / 502 Project “COSAVE: Regional Strengthening of the Implementation of Phytosanitary Measures and Market Access”.

The beneficiaries are COSAVE and the NPPOs of the seven countries that make up COSAVE. The Standards and Trade Development Facility (STDF) fund it, the Inter-American Institute for Cooperation on Agriculture (IICA) is the implementing organization and the International Plant Protection Convention (IPPC) Secretariat supports the project.

The editorial coordination was in charge of Maria de Lourdes Fonalleras and Florencia Sanz.

Maria de Lourdes Fonalleras, Florencia Sanz y Alec McClay, have defined the original structure of this Guide. The content development corresponds exclusively to Alec McClay expert contracted especially for the project.

The technical readers that made important contributions to the content of the case studies are the specialists of the NNPO's participating in the Project:

Adriana Ceriani, Melisa Nedilskyj, Leonardo Emilio Simón y Marcelo Sánchez from Servicio Nacional de Sanidad y Calidad Agroalimentaria – SENASA from Argentina;

Víctor Manuel Lima y Carla Roca Orellanos from Servicio Nacional de Sanidad Agropecuaria e Inocuidad Alimentaria – SENASAG from Bolivia;

Adriana Araújo Costa Truta y Clidenor Mendes Wolney Valente from Secretaria de Defesa Agropecuaria – SDA/MAPA from Brasil;

Cecilia Niccoli y Lilian Daisy Ibáñez from Servicio Agrícola y Ganadero – SAG from Chile;

Maria Eugenia Villalba y Mirta Zalazar from Servicio Nacional de Calidad, Sanidad Vegetal y de Semillas – SENAVER from Paraguay;

Efraín Arango Ccente y Cecilia Lévano Stella from Servicio Nacional de Sanidad Agraria – SENASA from Perú;

Leticia Casanova y María José Montelongo from Dirección General de Servicios Agrícolas – DGSA/ MGAP from Uruguay.

We express special appreciation to all of them. We also thank the support received from the IPPC Secretariat for the implementation of this component of the project.

Finally, we thanks Esteban Grille by diagramming the document.

**RISK ANALYSIS FOR
PLANTS AS PESTS
FOR *Ambrosia trifida* L.
(Asteraceae)**



Ambrosia trifida

Image: Theodore Webster, USDA Agricultural Research Service

TABLE OF CONTENTS

1. Stage I: Initiation	6
1.1. Initiation point for the pest risk analysis (PRA).....	6
1.2. Identity of the plant.....	6
1.3. Identification of the pest risk analysis area.....	7
1.4. Pest risk analysis history.....	7
2. Stage II. Weed risk assessment	8
2.1. Categorization.....	8
2.1.1. Presence or absence of the plant in the pest risk analysis area	
2.1.2. Regulatory status	
2.1.3. Potential for establishment and spread in the pest risk analysis area	
2.1.4. Potential for economic or environmental impact	
2.1.5. Conclusion of categorization	
2.2. Information about the plant.....	9
2.2.1. Geographic distribution of the plant	
2.2.2. Biology of the plant	
2.3. Risk Evaluation.....	13
2.3.1. Probability of introduction and spread	
2.3.2. Conclusion on the probability of introduction and spread	
2.3.3. Evaluation of potential economic and environmental consequences	
2.3.4. Conclusions on potential economic and environmental consequences	
2.4. Summary of the potential risk of <i>Ambrosia trifida</i>	18
3. Stage III: Pest risk management	19
4. References	20
Annex 1: Climatic tables.....	24
Annex 2: Method of combining probabilities and uncertainties.....	26

1. STAGE I: INITIATION

■ 1.1. INITIATION POINT FOR THE PEST RISK ANALYSIS

Ambrosia trifida was identified as a species which could be unintentionally introduced to the COSAVE region, as a contaminant of grains or other imported products. It is an important crop weed in North America and Europe, as well as being an important cause of allergies.

■ 1.2. IDENTITY OF THE PLANT

Accepted scientific name:

Ambrosia trifida L. (The Plant List, 2013)

Synonyms:

Ambrosia aptera DC, *Ambrosia integrifolia* Muhl. ex Willd. (The Plant List, 2013).

Common names:

In English: giant ragweed (official name according to the Weed Science Society of America), buffalo-weed, horseweed (Germplasm Resources Information Network, 2018), great ragweed (USDA-NRCS, 2018), Texan great ragweed, tall ragweed, blood ragweed, perennial ragweed (Integrated Taxonomic Information System, 2018).

In Spanish: artemisa grande (EPPO, 2018a), although no examples were found of the use of this name in the literature, and it is sometimes applied to *Artemisia tridentata* Nutt.

No Portuguese common name for *A. trifida* was found.

Taxonomic position:

Family Asteraceae

Subfamily Asteroideae

Tribe Heliantheae

(Funk et al., 2009).

Ambrosia trifida was described by Linnaeus in 1753 (IPNI, 2018). It is a well-defined species that presents no taxonomic or identification problems (Strother, 2006).

■ 1.3. IDENTIFICATION OF THE PEST RISK ANALYSIS AREA

For the purpose of this case study, the PRA area will be considered as the entire COSAVE region.

■ 1.4. PEST RISK ANALYSIS HISTORY

No previous pest risk analysis (PRA) performed by any COSAVE member country was found.

In Australia *A. trifida* was identified as a species that could be introduced as a contaminant of maize imported from the USA. (Weed Technical Working Group, 1999). A PRA of *Ambrosia artemisiifolia*, *Ambrosia psilostachya* and *Ambrosia trifida* for Poland concluded that these species warranted classification as quarantine pests (Karnkowski, 2001). However, they do not appear in the current list of species regulated by Poland (EPPO, 2018b). Mekky et al. (2010) analyzed the risk of introduction of *Ambrosia trifida*, *Ambrosia artemisiifolia* and other weeds in Egypt by the pathway of contamination of imported grains.

■ 1.5. CONCLUSION OF STAGE I

A pest risk analysis was carried out for *A. trifida* for the COSAVE region as a whole, in response to the risk of its unintentional introduction.

2. STAGE II. WEED RISK ASSESSMENT

■ 2.1. CATEGORIZATION

— 2.1.1. PRESENCE OR ABSENCE OF THE PLANT IN THE PEST RISK ANALYSIS AREA

No records of *A. trifida* were found for the COSAVE region in GBIF (Global Biodiversity Information Facility, 2018), Flora of Argentina (Zuloaga, 2006), Flora of Brazil (Flora do Brasil, 2020 em construção), or in the database of non-native plants in Chile (Fuentes et al., 2013). It was concluded that *A. trifida* is absent from the COSAVE region.

— 2.1.2. REGULATORY STATUS

— 2.1.2.1. In the pest risk analysis area

Ambrosia trifida does not appear in the List of Principal Regulated Pests for the COSAVE Region (COSAVE, 2016). In Peru it is classified as a quarantine pest not present (SENASA-PERÚ, 2017). *Ambrosia trifida* appears on the list of regulated pests for Argentina as a quarantine pest, not present (IPPC, 2017).

— 2.1.2.2. Worldwide

Ambrosia trifida appears on the A1 lists (absent quarantine pests) of Kazakhstan, Uzbekistan, Azerbaijan, and Ukraine; in the A2 lists (quarantine pests not widely distributed) of Moldova and Russia; and is a quarantine pest in Belarus. It also appears in the EPPO list of Invasive Alien Plants (EPPO, 2018a); EPPO strongly recommends that its member countries take measures to prevent the introduction and spread of the species mentioned in this list.

In South Africa *A. trifida* is a prohibited alien species under the National Environmental Management: Biodiversity Act (Department of Environmental Affairs, 2014).

In Canada, *A. trifida* seed is classified as “Class 2 primary noxious” under the Federal Seed Act, which limits the amount of weed seeds allowed in seeds offered for sale or imported into Canada (Canadian Food Inspection Agency, 2017).

In the U.S.A. *A. trifida* is classified as a state noxious weed by the states of California, Delaware and Illinois (USDA-NRCS, 2018).

In Egypt the importation of grains contaminated with seeds of *A. trifida*, *A. psilostachya* and *A. artemisiifolia* is prohibited, according to (Mekky et al., 2010). However, these species do not appear in the current list of pests regulated by Egypt, which in fact does not include any weed species (IPPC, 2014).

— 2.1.3. POTENTIAL FOR ESTABLISHMENT AND SPREAD IN THE PRA AREA

There are potentially suitable environmental and climatic conditions for *A. trifida* in parts of the territory of all COSAVE member countries (see 2.3.2).

— 2.1.4. POTENTIAL FOR ECONOMIC OR ENVIRONMENTAL IMPACT

Ambrosia trifida is a weed of great economic importance in corn, soybeans, wheat, cotton, and other crops in North America and China, and to a lesser extent in Europe. Its effects on human health due to the production of allergies also have a considerable economic impact (see 2.3.3.3).

— 2.1.5. CONCLUSION OF CATEGORIZATION

Based on the information gathered for the previous sections, it was concluded that *A. trifida* meets the requirements to be considered as a quarantine pest, being absent from the PRA area and with potential to cause economic or environmental impacts on plants in the PRA area.

■ 2.2. INFORMATION ABOUT THE PLANT

— 2.2.1. GEOGRAPHIC DISTRIBUTION OF THE PLANT

Native distribution:

Canada: Alberta, Manitoba, New Brunswick, Nova Scotia, Ontario, Prince Edward Island, Quebec, Saskatchewan (USDA-NRCS, 2018).

USA: Alabama, Arizona, Arkansas, California, North Carolina, South Carolina, Colorado, Connecticut, North Dakota, South Dakota, Delaware, District of Columbia, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New York, New Mexico, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, Tennessee, Texas, Utah, Vermont, Virginia, West Virginia, Washington, Wisconsin, Wyoming (USDA-NRCS, 2018).

Mexico: Chihuahua, Coahuila, Sonora, Baja California Sur (Germplasm Resources Information Network, 2018; Global Biodiversity Information Facility, 2018)

Naturalized distribution:

Asia: Georgia, China, Japan (Germplasm Resources Information Network, 2018); Korea (Kim et al., 2004). In China it is reported from the provinces of Hebei, Heilongjiang, Hunan, Jiangxi, Jilin, Liaoning, Shandong, Sichuan, and Zhejiang (Flora of China Editorial Committee, 2011).

Europe: Lithuania, Germany, the Netherlands, Slovakia, Denmark, United Kingdom, Italy, France, Spain (Germplasm Resources Information Network, 2018); Austria, Slovenia, Switzerland, Czech Republic, Serbia (Follak et al., 2013).

In addition to the countries mentioned where *A. trifida* is naturalized, there are casual records or records of unknown status of *A. trifida* in Israel, Belarus, Denmark, Estonia, Finland, Latvia, Moldova, Norway, Russia (European part), Sweden, Ukraine, Belgium, Poland, Ireland, and Slovenia (DAISIE, 2018; Germplasm Resources Information Network, 2018).

Cultivated distribution:

Ambrosia trifida is not cultivated.



Figure 1. Worldwide distribution of *Ambrosia trifida* (Global Biodiversity Information Facility, 2018); provinces of China with records of *A. trifida* according to Flora of China Editorial Committee (2011).

— 2.2.2. BIOLOGY OF THE PLANT

— 2.2.2.1. Morphology

Ambrosia trifida is a herb with branched, erect stems 30 - 150 cm in height (sometimes up to 400 cm). Leaves mostly opposite, slightly scabrous, rounded-deltate to ovate or elliptical, 40-150 × 30-70 mm, some palmate with 3 (-5) lobes, margins usually dentate, petioles 10-30 mm. Flowers green, monoecious, inflorescence in the form of a spike. The fruit is a pyramidal achene 3-5 mm in length, with 4 to 5 straight spines of 0.5-1 mm, containing a single seed (Strother, 2006).

— 2.2.2.2. Life cycle

Ambrosia trifida is annual and blooms from July to November in the northern hemisphere. Pollination is by wind (anemophilous). It reproduces only by seed. The flowers are capable of self-pollination, but the progeny from this are less vigorous than those resulting from crossing. A typical *A. trifida* plant in Illinois produces about 275 seeds (Bassett and Crompton, 1982), although a production of 1,650 seeds per plant is also cited (Stevens, 1932). The mature seeds survive the winter persisting in the inflorescences or in the soil.

In Illinois (USA) the seeds of *A. trifida* are among the first species to germinate in the spring, at the beginning of March. In Quebec (Canada) the time of maximum germination is from the end of April to the beginning of May. In the U.S.A. some populations of *A. trifida* show multiple flushes of germination throughout the growing season (Michigan State University, 2018). Seeds germinate best at alternating temperatures of 20°C - 30°C, after a cold period (stratification), and covered by a minimum of 2 cm of soil (Bassett and Crompton, 1982).

The viability of freshly collected seeds in Ohio (USA) varied between 48% and 53%. In a period of 4 years, the percentage of viable seeds dropped to zero for the seeds on the surface of the soil and 19% when they were buried at a depth of 20 cm; however, some seeds buried at 20 cm can retain their viability at least 9 years in the soil (Harrison et al., 2007).

— 2.2.2.3. Dispersal

The seeds of *A. trifida* have no specific adaptations for dispersal. The seeds do not float well, indicating that dispersal by water is not very important (Parker and Leck, 1985). However, seeds of *A. trifida* were found in very small amounts among leaf litter deposited on marine beaches and along the River Rhine in the Netherlands (Cappers, 1993).

In Ohio (USA) the European earthworm *Lumbricus terrestris* collects the seeds of *A. trifida* and buries them in its burrows at a depth of up to 22 cm, removing a large amount of the seed that falls on the soil surface. This protects the seed from other predators and may contribute to the formation of the seed bank (Regnier et al., 2008).

— 2.2.2.4. Habitat and environmental factors affecting the plant

Ambrosia trifida is a plant of alluvial plains, and predominates in disturbed, moist soils along irrigation canals and streams (Bassett and Crompton, 1982). Strother, (2006) indicates its habitat as disturbed sites and vacant lots with wet soils. In New Jersey (USA) it was abundant on the banks of channels in a freshwater tidal wetland (Parker and Leck, 1985).

In central and eastern Europe it is mainly a species of ruderal habitats (such as industrial and urban areas, and railroad edges), and is found less frequently in riparian zones or cultivated fields (Follak et al., 2013). In South Korea it was abundant in a disused landfill near Seoul (Kim et al., 2004) and in a riverbank area (Lee et al., 2010). In Japan it is present in farms, orchards, paddocks, riverbanks, roadsides, and vacant lots (National Research and Development Agency, 2018) and was found invading a nature reserve (Miyawaki and Washitani, 1996, cited in Follak et al. al., 2013).

Ambrosia trifida has no specific requirements as to the type of soil (CABI, 2016) but according to the University of Michigan (2018) prefers fertile soils with high levels of nitrogen, phosphorus and potassium

— 2.2.2.5. Climatic Adaptation

The great majority of records of *A. trifida* with geographical coordinates in GBIF occur in climatic zones Dfb (snow climate, fully humid, warm summer), Dfa (continental without dry season, warm summer, cold winter) [not present in the COSAVE region], Cfa (warm temperate, fully humid, hot summer), Cfb (warm temperate, fully humid, warm summer) and BSk (cold steppe) according to the modified Köppen and Geiger system (see Annex 1), with a few localities in zones Csb, Csa, Dfc, BSh, BWh, and Dwa (Figure 2).




Figure 2. Worldwide distribution of *Ambrosia trifida* in relation to the modified Köppen-Geiger climate system.

In terms of the NAPPFAST system (Magarey et al., 2008), the great majority of the world distribution of *A. trifida* is found in zones 4 to 9, with a few locations in zone 3 in Canada, (Figure 3) corresponding to minimum annual temperature of -40°C to -9.4°C (see Table A2 in Annex). Since *A. trifida* is an annual species that survives the winter in the seed state, it is possible that its distribution is limited in cold climates not so much by low winter temperatures but by the lack of heat during the growing season. The absence of *A. trifida* from zones 10 to 13 may be related to the need for a cold period to break dormancy of the seeds (Davis et al., 2013).




Figure 3. Worldwide distribution of *Ambrosia trifida* in relation to the NAPPFAST classification of cold hardiness zones.

In central and eastern Europe, *A. trifida* is associated with regions of more marked seasonality of precipitation (Follak et al., 2013).

— 2.2.2.6. Methods of control

Ambrosia trifida is considered one of the most difficult weed species to control with herbicides due to its rapid growth, the occurrence of multiple flushes of germination, and the ability of the seeds to emerge from a depth of up to 15 cm where the applied herbicides do not penetrate the soil (Michigan State University, 2018).

There are many recommendations for control programs of *A. trifida* with herbicides, depending on the crop and production system (Johnson et al., 2007; United Soybean Board, 2016).

The control of *A. trifida* in the USA and Canada is complicated by the existence of many herbicide-resistant populations, including to Group 2 products (chlorimuron-ethyl, cloransulam-methyl, imazamox, imazaquin, imazethapyr, primisulfuron-methyl, prosulfuron) and Group 9 products (glyphosate). These populations include several resistant to multiple herbicides, including cases of combined resistance to Group 2 and Group 9 (Heap, 2018).

Several North American insect species have been introduced in Europe, China and Australia as biological control agents for *A. artemisiifolia* (Gerber et al., 2011). It is possible that some of these also attack *A. trifida* since they are generally genus-specific. An American beetle, *Ophraella communa*, accidentally introduced in Europe, seems to have a significant impact on the populations of *A. artemisiifolia* (Müller-Schärer et al., 2014); its potential impact on *A. trifida* is unknown.

■ 2.3. RISK EVALUATION

— 2.3.1. PROBABILITY OF INTRODUCTION AND SPREAD

— 2.3.1.1. Probability of entry

Natural dispersal

Given the distances between the COSAVE region and the areas where *A. trifida* is present, and the lack of long-distance dispersal mechanisms in this species, natural dispersal does not appear to be an important entry pathway to the region.

Unintentional introduction

In Europe *A. trifida* was introduced as a contaminant of cereal grains and oilseeds imported from North America. Specifically, it arrived in Germany with imports of spring wheat seed before 1906, and Slovakia with grains imported from North America via the Soviet Union (Follak et al., 2013); these authors suggest that the frequency of introduction of *A. trifida* in Europe has decreased in recent times due to improvements in seed cleaning methods.

In Belgium (Verloove, 2006) indicates that *A. trifida* was introduced with imported grains and wool, without further details of the date of introduction or the source of the information.

(Mekky et al., 2010) reports that in grain of wheat, corn and sorghum imported into Egypt between 2009 and 2010, contamination with seeds of *Ambrosia* spp. (including *A. trifida*) was found in 3.7% of shipments originating in Ukraine, 1% in the USA,

and 2.3% in Russia. The maximum level of contamination was 144 seeds per kg for Ukraine, 2 seeds per kg for the USA. and 4 seeds per kg for Russia.

In Australia, between 1994 and 1995, seeds of *A. trifida* were detected in imports of corn and sorghum from the USA and of soybean (origin not indicated) (Weed Technical Working Group, 1999).

(Karnkowski, 2001) reports many interceptions of *A. trifida* and other *Ambrosia* species in Russia, Finland and Poland, summarized in Table 1.

In Peru, *A. trifida* was detected on 43 occasions in shipments of corn and soybean grain imported from the USA in 2017 and 2018. In the period 2008 - 2016 there were no records of interception of *Ambrosia* (SENASA-PERÚ, 2018).

Intentional introduction

No evidence of pathways for intentional introduction was found.

Table 1. Interceptions of *A. trifida* and other *Ambrosia* species in products imported into some European countries, according to (Karnkowski, 2001).

Species	Product	Country(s) or region of origin	Importing country
<i>Ambrosia trifida</i>	Corn and sorghum grain	Canada, USA	Russia
	Soybean grain	USA	Russia
	Grass seed	Netherlands	Russia
	Rice grain	Japan	Russia
	Barley grain	France, Canada	Russia
	Soy meal	Germany, Netherlands, Brazil	Russia
	Corn, soybean and rye grain	North America	Finland
<i>Ambrosia</i> spp.	Corn and soybean grain, soy meal	USA	Poland
	Soybean grain	Canada	Poland
	Corn, wheat, triticale, soybean, millet, sunflower grain and corn meal	Hungary	Poland
	Corn, wheat, barley and sunflower grain	Slovakia	Poland
	Corn, sunflower and millet grain	Czech Republic	Poland
	Corn, wheat, millet, white mustard, buckwheat and sunflower grain and medicinal herbs	Ukraine	Poland
	Sunflower grain	Belarus	Poland
	Soybean grain	Netherlands, Romania	
	Corn and soybean grain	Austria	Poland
	Corn grain	France	Poland
	Sunflower and buckwheat grain	Russia	Poland
	Soybean meal	Germany	Poland
	Sunflower grain	Belarus	Poland

The possible pathways of entry for *A. trifida* are summarized in Table 2.

Table 2. Potential entry pathways for *Ambrosia trifida*

Entry pathway	Probability	Uncertainty
1. Contaminant of seed imported for sowing	medium	medium
2. Contaminant of imported grain	low	medium
3. Contaminant of imported wool	negligible	negligible

2.3.1.2. Probability of establishment

The percentage of the territory of the member countries of COSAVE included in the most favorable climatic zones for *A. trifida* (BSk, Cfa, Cfb, Dfb) varies between 7.7% for Brazil up to 100% for Uruguay. If zones BSh, BWh, Csb, and Dfc, which are also within the climatic range of *A. trifida*, are added, these percentages increase slightly (Table 3). In the COSAVE region, the areas of zone NAPPFAST 1 and 2, outside the cold limit for *A. trifida*, are insignificant (Table A2 in Annex), indicating that this species would not be limited by low temperatures in any part of the region. In contrast, significant areas of Bolivia, Brazil and Peru are in zones 11 to 13, and probably have climates which are too tropical for *A. trifida* (Table 3).

Within the climatically favorable areas for *A. trifida*, habitats potentially suitable for establishment would include annual crop fields, disturbed and urban areas, vacant lots, roads and railways, wetlands and riparian areas. These habitats exist in all member countries of COSAVE

Table 3. Percentage of the territory of each COSAVE member country included in the indicated Köppen-Geiger and NAPPFAST climate zones (see Annex 1, Tables A1 and A2).

Country	Köppen-Geiger zones		NAPPFAST Zones
	BSk, Cfa, Cfb, Dfb	BSh, BSk, BWh, Cfa, Cfb, Csb, Dfb, Dfc	3 - 10
Argentina	53.1%	68.0%	100.0%
Bolivia	11.3%	18.0%	49.4%
Brazil	7.7%	13.5%	12.2%
Chile	14.3%	33.1%	81.7%
Paraguay	36.2%	54.5%	99.9%
Peru	8.4%	17.8%	32.4%
Uruguay	100.0%	100.0%	99.0%

Based on the above, the probability of establishment of *A. trifida* is rated as high with low uncertainty, considering the COSAVE region as a whole.

— 2.3.1.3. Probability of spread

Natural dispersal

In Europe the dispersion of *A. trifida* has been relatively slow and it does not tend to colonize areas outside its main habitat, which are ruderal zones and railroads, perhaps due to its low fecundity and low seed viability (Follak et al., 2013)

Unintentional dispersal

No specific data were found on the unintentional dispersion of *A. trifida*. It can be assumed that the seed can be transported as a contaminant of agricultural products such as grains and seeds from infested areas, and also by the movement of vehicles, agricultural machinery, or animals.

Intentional dispersal

Ambrosia trifida has no uses in agriculture, horticulture or as a medicinal plant. Therefore, there seems to be no motive for intentional dispersal.

Potential pathways of spread for *A. trifida* are summarized in Table 4.

Table 4. Potential pathways of spread for *Ambrosia trifida* within the COSAVE region

Pathway of spread	Probability	Uncertainty
Natural dispersal by water or wind	low	low
Contaminant of agricultural products	medium	medium
Unintentional transportation with vehicles, agricultural machinery, or animals	medium	medium
Intentional dispersal, for example for seeding	negligible	low
Overall probability of spread	medium	medium

Based on the above, the probability of spread of *A. trifida* is rated as medium with medium uncertainty.

— 2.3.2. CONCLUSION ON THE PROBABILITY OF INTRODUCTION AND SPREAD

Combining the probabilities according to the method of Annex 2, it is concluded that the overall probability of introduction and spread of *A. trifida* is rated as medium with medium uncertainty.

— 2.3.3. EVALUATION OF POTENTIAL ECONOMIC AND ENVIRONMENTAL CONSEQUENCES

— 2.3.3.1. Economic effects

Effects on crop yield or quality

Ambrosia trifida is an important crop weed in North America, especially in soybean (*Glycine max*), sunflower (*Helianthus annuus*), bean (*Phaseolus spp.*), corn (*Zea mays*),

wheat (*Triticum* spp.) and cotton (*Gossypium hirsutum*) (CABI, 2016). There is an extensive literature on its impacts and control. In experiments carried out in Missouri, USA, for two years, dense *A. trifida* populations reduced soybean seed yields by approximately 50% (Baysinger and Sims, 1991). There was also a 55% reduction in corn yield in Michigan (Michigan State University, 2018). A density of 1 plant of *A. trifida* per m² reduced the yield of sweet corn by approximately 40% and affected several parameters of crop quality (Williams and Masiunas, 2006). In Tennessee a density of 0.26 plants of *A. trifida* per metre of row reduced the yield of cotton by 50% (Barnett and Steckel, 2013).

In Northeast China *A. trifida* is considered one of the weeds that causes the most economic damage to wheat and other annual crops. It was found that the plant and its residues have allelopathic effects that reduce wheat growth (Kong et al., 2007).

All crops affected by *A. trifida* as weeds in the areas where it is currently present are of economic importance for the COSAVE region.

Effects on production costs

No specific and recent data on the effects of *A. trifida* on production costs were found. However, the need for herbicide application would probably entail additional costs for producers in the event that *A. trifida* becomes established in the region.

Commercial effects

Given that *Ambrosia trifida* is a prohibited or quarantine pest in several countries (see 2.1.2.), its presence in a COSAVE member country could have an impact on market access for exported products such as grains or seeds.

Social effects

The pollen of *Ambrosia* spp. (including *A. trifida*) is an important cause of allergies that produce serious suffering in affected populations and areas (see 2.4.3).

— 2.3.3.2. Environmental effects

Effects on plant species

In general, *A. trifida* is a species that colonizes disturbed and cultivated areas that do not tend to have great importance for biodiversity (Plank et al., 2016), and therefore, few impacts on native plant species or communities have been identified in areas where *A. trifida* has invaded. However, in Japan the diversity of plant species had a negative correlation with abundance of *A. trifida* in a riparian natural reserve near Tokyo (Miyawaki and Washitani, 1996), and unspecified impacts on native plants such as *Primula sieboldii* E. Morren (Primulaceae) are reported. (National Research and Development Agency, 2018).

Effects on ecological systems or processes

No evidence was found of effects of *A. trifida* on ecological systems or processes.

— 2.3.3.3. Non-phytosanitary effects

Species of *Ambrosia* (including *A. trifida* and *A. artemisiifolia*) are among the most important causes of respiratory allergies (hay fever) in North America and Europe, due to their abundant production of wind-dispersed pollen. In Germany it was calculated in 2003 that the annual economic cost of allergies caused by *A. artemisiifolia* was between €20 - €50 million (USD25 - 62 millions), and in a single hospital in Italy the annual cost for allergy treatment of *Ambrosia* was € 1.3 million

(USD \$1.63 millions) (Plank et al., 2016). With climate change and the increase of atmospheric CO₂ it is expected that pollen production by *Ambrosia* spp. will increase (Rogers et al., 2006).

— 2.3.4. CONCLUSIONS ON POTENTIAL ECONOMIC AND ENVIRONMENTAL CONSEQUENCES

Summing up all the potential consequences identified, they are rated as high with low uncertainty.

■ 2.4. SUMMARY OF THE POTENTIAL RISK OF *AMBROSIA TRIFIDA*

Ambrosia trifida is a species of great importance as a weed of crops, difficult to control and causing significant yield reductions in several crops. There are suitable conditions for its establishment in all the countries of the region. The entry pathway that presents the highest degree of risk to the region is the importation of contaminated seed for planting. The potential risk of *A. trifida* for the COSAVE region is summarized in Table 5.

Table 5. Summary of the potential risk of *Ambrosia trifida* for the COSAVE region.

	Risk rating	Uncertainty
Probabilities of entry		
1. Contaminant of seed imported for sowing	Medium	Medium
2. Contaminant of imported grain	Low	Medium
3. Contaminant of imported wool	Negligible	Negligible
Probability of establishment	High	Low
Probability of spread	Medium	Medium
Overall probability of establishment and spread	Medium	Medium
Consequences		
Potential economic and environmental consequences	High	Low

3. STAGE III: PEST RISK MANAGEMENT

Based on the foregoing, it is recommended that *Ambrosia trifida* be included in the list of quarantine pests, and that the following phytosanitary measures be applied:

Requirements for imported seed for sowing:

DA¹ 5. The place of production / site of production site / field, was inspected during the growing season and found to be free of *Ambrosia trifida*,

or

DA 15. The shipment is free of *Ambrosia trifida*, according to the result of an official laboratory analysis.

1 Additional Declaration (DA) statement that is required by an importing country to be entered on a phytosanitary certificate and which provides specific additional information on a consignment in relation to regulated pests or regulated articles (ISPM n°5)

4. REFERENCES

- Barnett, K.A. and L.E. Steckel. 2013. Giant ragweed (*Ambrosia trifida*) competition in cotton. *Weed Science* 61: 543-548.
- Bassett, I.J. and C.W. Crompton. 1982. The biology of Canadian weeds. 55. *Ambrosia trifida* L. *Canadian Journal of Plant Science* 62: 1003-1010.
- Baysinger, J.A. and B.D. Sims. 1991. Giant ragweed (*Ambrosia trifida*) Interference in soybeans (*Glycine max*). *Weed Science* 39: 358-362.
- CABI. 2016. *Invasive Species Compendium: Ambrosia trifida (giant ragweed)*. Accessed online February 13 2018. <https://www.cabi.org/isc/datasheet/4693>.
- Canadian Food Inspection Agency. 2017. *Weed Seed: Ambrosia trifida (Giant ragweed)*. Accessed online February 13 2018. <http://www.inspection.gc.ca/plants/seeds/testing-grading/seeds-identification/ambrosia-trifida/eng/1472605410597/1472605411018>.
- Cappers, R.T.J. 1993. Seed dispersal by water: a contribution to the interpretation of seed assemblages. *Vegetation History and Archaeobotany* 2: 173-186.
- COSAVE. 2016. *Listado de las Principales Plagas Reglamentadas para la Región del COSAVE*. Accessed online January 29 2018. <http://www.cosave.org/sites/default/files/paginas/adjuntos/Anexo%20Resol%20213%20%20principales%20plagas%20reglamentadas.pdf>.
- DAISIE. 2018. *Species Factsheet: Ambrosia trifida*. Accessed online February 13 2018. <http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21722#>.
- Davis, A.S., S. Clay, J. Cardina, A. Dille, F. Forcella, J. Lindquist and C. Sprague. 2013. Seed burial physical environment explains departures from regional hydrothermal model of giant ragweed (*Ambrosia trifida*) seedling emergence in U.S. Midwest. *Weed Science* 61: 415-421.
- Department of Environmental Affairs. 2014. National List of Invasive Species in Terms Section 70(1)(A). *Government Gazette (South Africa)* 37886: 8-31.
- EPPO. 2018a. *EPPO Global Database: Ambrosia trifida (AMBTR)*. Accessed online February 13 2018. <https://gd.eppo.int/taxon/AMBTR>.
- EPPO. 2018b. *EPPO Global Database: Regulated organisms in Poland*. Accessed online February 13 2018. <https://gd.eppo.int/country/PL/regulated>.
- Flora do Brasil. 2020 em construção. Accessed online February 10 2018. <http://floradobrasil.jbrj.gov.br/>.
- Flora of China Editorial Committee. 2011. *Ambrosia trifida Linnaeus*. Accessed online February 13 2018. http://www.efloras.org/florataxon.aspx?flora_id=2&taxon_id=200023073.
- Follak, S., S. Dullinger, I. Kleinbauer, D. Moser and F. Essl. 2013. Invasion dynamics of three allergenic invasive Asteraceae (*Ambrosia trifida*, *Artemisia annua*, *Iva xanthiifolia*) in central and eastern Europe. *Preslia* 85: 41-61.

- Fuentes, N., A. Pauchard, P. Sánchez, J. Esquivel and A. Marticorena. 2013. A new comprehensive database of alien plant species in Chile based on herbarium records. *Biological Invasions* 15: 847-858.
- Funk, V.A., T. Stuessy and R. Bayer. 2009. Systematics, Evolution, and Biogeography of Compositae, International Association for Plant Taxonomy Vienna, Austria.
- Gerber, E., U. Schaffner, A. Gassmann, H.L. Hinz, M. Seier and H. Müller-Schärer. 2011. Prospects for biological control of *Ambrosia artemisiifolia* in Europe: learning from the past. *Weed Research* 51: 559-573.
- Germplasm Resources Information Network. 2018. *Ambrosia trifida* L. Accessed online February 10 2018. <https://npgsweb.ars-grin.gov/gringlobal/taxonomydetail.aspx?id=103827>.
- Global Biodiversity Information Facility. 2018. *Ambrosia trifida* L. Accessed online January 29 2018. <https://www.gbif.org/species/3110588>.
- Harrison, S.K., E.E. Regnier, J.T. Schmoll and J.M. Harrison. 2007. Seed size and burial effects on giant ragweed (*Ambrosia trifida*) emergence and seed demise. *Weed Science* 55: 16-22.
- Heap, I. 2018. *The International Survey of Herbicide Resistant Weeds*. Accessed online February 14 2018. <http://www.weedscience.org/Summary/Species.aspx?WeedID=184>.
- Integrated Taxonomic Information System. 2018. *Ambrosia trifida* L. Accessed online February 11 2018. https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=36521#null.
- IPNI. 2018. *International Plant Names Index: Ambrosia trifida* L. Accessed online February 12 2018. <http://www.ipni.org/ipni/idPlantNameSearch.do?id=315739-2>.
- IPPC. 2014. *List of regulated pests [Egypt]*. Accessed online February 14 2018. <https://www.ippc.int/en/countries/egypt/reportingobligation/2014/08/list-of-regulated-pests-3/>.
- IPPC. 2017. *Listado de Plagas Reglamentadas para Argentina*. Accessed online August 9 2018. <https://www.ippc.int/en/countries/argentina/reportingobligation/2017/09/quarantine-pest-list-of-argentina-2017/>.
- Johnson, B., M. Loux, D. Nordby, C. Sprague, G. Nice, A. Westhoven and J. Stachler. 2007. *Biology and Management of Giant Ragweed*. Accessed online February 14 2018. <https://www.extension.purdue.edu/extmedia/BP/GWC-12.pdf>.
- Karnkowski, W. 2001. *Pest Risk Analysis and Pest Risk Assessment for the territory of the Republic of Poland (as PRA area) on Ambrosia spp. (updated version)*. Accessed online February 14 2018. https://www.eppo.int/QUARANTINE/Pest_Risk_Analysis/PRAdocs_plants/08-14124%20PRA-Ambrosia.doc.
- Kim, K.D., E.J. Lee and K.-H. Cho. 2004. The plant community of Nanjido, a representative nonsanitary landfill in South Korea: implications for restoration alternatives. *Water, Air, & Soil Pollution* 154: 167-185.
- Kong, C.-H., P. Wang and X.-H. Xu. 2007. Allelopathic interference of *Ambrosia trifida* with wheat (*Triticum aestivum*). *Agriculture, Ecosystems & Environment* 119: 416-420.
- Lee, C.S., Y.C. Cho, H.C. Shin, G.S. Kim and J.H. Pi. 2010. Control of an invasive alien species, *Ambrosia trifida* with restoration by introducing willows as a typical riparian vegetation. *Journal of Ecology and Environment* 33: 157-164.

- Magarey, R.D., D.M. Borchert and J.W. Schlegel. 2008. Global plant hardiness zones for phytosanitary risk analysis. *Scientia Agricola* 65: 54-59.
- Mekky, M.S., E.E. Hassanein, A.N.M. Nassar, M.R. Moshtohry, A.S. Kholosy and M.F.I. Daie. 2010. Weed risk analysis and assessment of weed seed consignment with imported grains. *Egyptian Journal of Agricultural Research* 88.
- Michigan State University. 2018. *Giant Ragweed (Ambrosia trifida L.)*. Accessed online February 13 2018. <https://www.canr.msu.edu/weeds/extension/giant-ragweed>.
- Miyawaki, S. and I. Washitani. 1996. A population dynamics model for soil seedbank plant and its application to the prediction of the effects of weeding on a population of *Ambrosia trifida* L. invading into a nature reserve. *Japanese Journal of Conservation Ecology* 1: 25-47.
- Müller-Schärer, H., S.T.E. Lommen, M. Rossinelli, M. Bonini, M. Boriani, G. Bosio and U. Schaffner. 2014. *Ophraella communa*, the ragweed leaf beetle, has successfully landed in Europe: fortunate coincidence or threat? *Weed Research* 54: 109-119.
- National Research and Development Agency. 2018. *Invasive Species of Japan: Ambrosia trifida*. Accessed online February 15 2018. <http://www.nies.go.jp/biodiversity/invasive/DB/detail/80410e.html>.
- Parker, V.T. and M.A. Leck. 1985. Relationships of seed banks to plant distribution patterns in a freshwater tidal wetland. *American Journal of Botany*: 161-174.
- Plank, L., D. Zak, M. Getzner, S. Follak, F. Essl, S. Dullinger, I. Kleinbauer, D. Moser and A. Gattringer. 2016. Benefits and costs of controlling three allergenic alien species under climate change and dispersal scenarios in Central Europe. *Environmental Science & Policy* 56: 9-21.
- Regnier, E., S.K. Harrison, J. Liu, J.T. Schmoll, C.A. Edwards, N. Arancon and C. Holloman. 2008. Impact of an exotic earthworm on seed dispersal of an indigenous US weed. *Journal of Applied Ecology* 45: 1621-1629.
- Rogers, C.A., P.M. Wayne, E.A. Macklin, M.L. Muilenberg, C.J. Wagner, P.R. Epstein and F.A. Bazzaz. 2006. Interaction of the onset of spring and elevated atmospheric CO₂ on ragweed (*Ambrosia artemisiifolia* L.) pollen production. *Environmental Health Perspectives* 114: 865-869.
- SENASA-PERÚ. 2017. Servicio Nacional de Sanidad Agraria: Lista de Plagas Cuarentenarias no Presentes en el Perú.
- SENASA-PERÚ. 2018. Servicio Nacional de Sanidad Agraria: Reporte de Sistema Integrado de Gestión de Sanidad Vegetal -SIGSVE.
- Stevens, O.A. 1932. The number and weight of seeds produced by weeds. *American Journal of Botany* 19: 784-794.
- Strother, J.L. 2006. *Ambrosia* Linnaeus, pp. 10-18. In *Flora of North America* Editorial Committee (ed.), *Flora of North America: North of Mexico* vol 21 Magnoliophyta: Asteridae, Part 8: Asteraceae, Part 3. Oxford University Press.
- The Plant List. 2013. *Ambrosia trifida* L. Accessed online February 15 2018. <http://www.theplantlist.org/tpl1.1/record/gcc-7636>.
- United Soybean Board. 2016. *Management of Herbicide-Resistant Giant Ragweed*. Accessed online February 14 2018. https://weeds.cscience.missouri.edu/publications/FactSheet_GiantRagweed.pdf.

- USDA-NRCS. 2018. *The PLANTS Database: Ambrosia trifida L.: great ragweed*. Accessed online February 10 2018. <https://plants.usda.gov/core/profile?symbol=AMTR>.
- Verloove, F. 2006. Catalogue of neophytes in Belgium. *Scripta Botanica Belgica* 39: 1-89.
- Weed Technical Working Group. 1999. *Weed Risk Analysis of a Proposed Importation of Bulk Maize (Zea mays) from the USA*. Accessed online February 14 2018. http://www.agriculture.gov.au/SiteCollectionDocuments/ba/memos/1999/plant/TWGP_4.doc.
- Williams, M.M. and J.B. Masiunas. 2006. Functional relationships between giant ragweed (*Ambrosia trifida*) interference and sweet corn yield and ear traits. *Weed Science* 54: 948-953.
- Zuloaga, F.O. 2006. *Flora Argentina: Plantas Vasculares de la Republica Argentina*. Accessed online February 10 2018. <http://www.floraargentina.edu.ar/>.

ANNEX 1: CLIMATIC TABLES

Table A1. Percentage of the territory of each COSAVE member country corresponding to each of the climatic zones of the Köppen-Geiger system. Calculated using the March 2017 updated version with data from 1986-2010 and with a resolution of 5 minutes, according to Kottek and Rubel².

		Country						
		Argentina	Bolivia	Brazil	Chile	Paraguay	Peru	Uruguay
Af	Equatorial rainforest, fully humid		2.24	16.07		0.69	41.38	
Am	Equatorial monsoon		13.39	20.48		4.94	9.62	
As	Equatorial savannah with dry summer			2.56				
Aw	Equatorial savannah with dry winter		46.43	46.06		37.00	4.98	
BSh	Steppe climate, hot	7.13	6.62	5.76		18.26	1.67	
BSk	Steppe climate, cold	25.02	8.98		3.05		1.95	
BWh	Desert climate, hot	2.08	0.02	<0.01	0.67		7.73	
BWk	Desert climate, cold	6.06	5.52		25.52		4.08	
Cfa	Warm temperate, fully humid, hot summer	23.76	0.52	6.89		36.21		99.17
Cfb	Warm temperate, fully humid, warm summer	4.36	1.85	0.82	11.23		6.48	0.83
Cfc	Warm temperate, fully humid, cool summer and cold winter	1.22	0.05		12.65		0.18	
Csb	Warm temperate with dry, warm summer and cold winter	5.67			18.11			
Csc	Warm temperate with dry, cool summer and cold winter	0.74			1.07			
Cwa	Warm temperate with dry winter, hot summer	15.85	2.51	1.15		2.90		
Cwb	Warm temperate with dry winter, warm summer	1.98	6.01	0.21			4.66	
Cwc	Warm temperate with dry winter, cool summer and cold winter	0.45	0.60				0.73	
Dfb	Snow climate, fully humid, warm summer	<0.01						
Dfc	Snow climate, fully humid, cool summer and cold winter	0.02						
Dsc	Snow climate with dry, cool summer and cold winter	0.07			0.02			
Dwb	Snow climate with dry winter, warm summer	0.01						
Dwc	Snow climate with cool summer and cold, dry winter	0.02						
EF	Polar climate	0.01			0.02		0.01	
ET	Tundra climate	5.55	5.25		27.64		16.51	

² Kottek, M. and F. Rubel. 2017. World Maps of Köppen-Geiger Climate Classification. Accessed online January 10 2018. <http://kooppen-geiger.vu-wien.ac.at/present.htm>

Table A2. Percentage of the territory of each COSAVE member country corresponding to each of the NAPPFAST cold hardiness zones³.

NAPPFAST Zone	Mean annual extreme minimum temperature (°C)	Country						
		Argentina	Bolivia	Brazil	Chile	Paraguay	Peru	Uruguay
1	< -45.6	<0.01	0.00	0.00	0.00	0.00	0.00	0.00
2	-45.9 — -40.0	0.01	0.00	0.00	0.02	0.00	0.00	0.00
3	-40.0 — -34.4	0.07	0.00	0.00	0.01	0.00	0.00	0.00
4	-34.4 — -28.9	0.67	0.00	0.00	0.25	0.00	0.00	0.00
5	-28.9 — -23.3	2.09	0.15	0.00	1.07	0.00	0.00	0.00
6	-23.3 — -17.8	4.22	1.70	0.00	4.78	0.00	0.93	0.00
7	-17.8 — -12.2	7.45	9.74	0.00	11.47	0.00	5.31	0.00
8	-12.2 — -6.7	17.25	12.07	0.07	16.17	0.00	7.87	0.00
9	-6.7 — -1.1	46.69	10.64	3.69	26.29	4.46	9.35	80.22
10	-1.1 — 4.4	21.55	15.11	8.43	21.67	95.46	8.91	18.74
11	4.4 — 10.0	0.00	38.51	18.52	14.30	0.08	19.59	1.03
12	10.0 — 15.6	0.00	12.08	44.55	3.93	0.00	42.89	0.00
13	> 15.6	0.00	0.00	24.73	0.04	0.00	5.14	0.00

³ Calculated with data courtesy of Dr. R. Magarey, see Magarey, R.D., D.M. Borchert and J.W. Schlegel. 2008. Global plant hardiness zones for phytosanitary risk analysis. *Scientia Agricola* 65: 54-59.

ANNEX 2: METHOD OF COMBINING PROBABILITIES AND UNCERTAINTIES

To rate the overall risk of establishment and spread, each probability is converted into a numerical score (negligible = 0, low = 1, medium = 2, high = 3), and the numerical scores are multiplied as follows:

$$\begin{aligned} & \textit{Probability of establishment and spread} \\ & = \\ & \textit{Probability of establishment} \\ & \times \\ & \textit{Probability of spread} \end{aligned}$$

This product is used to rate the overall probability of introduction and spread as follows:

Product (probability of establishment × probability of spread)	Overall rating for probability of establishment and spread
0	Negligible
1 – 3	Low
4 – 6	Medium
>6	High

Similarly, the uncertainty levels of the probabilities of establishment and spread are combined to arrive at an uncertainty score for the overall probability of establishment and spread. As before, the levels of uncertainty are converted into numerical scores (negligible = 0, low = 1, medium = 2, high = 3). Unlike the probabilities, the uncertainties are added:

$$\begin{aligned} & \textit{Uncertainty of the probability of establishment and spread} \\ & = \\ & \textit{Uncertainty of the probability of establishment} \\ & + \\ & \textit{Uncertainty of the probability of spread} \end{aligned}$$

This sum is used to rate the uncertainty of the overall probability of establishment and spread as follows:

Sum of uncertainty scores for the overall probability of establishment and spread	Overall uncertainty rating for the probability of establishment and spread
0	Negligible
1	Low
2 – 3	Medium
4 – 6	High